



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

These are complete elliptic integrals of the second and first kinds respectively, and can be readily evaluated. See Peirce's *Short Table of Integrals*, pp. 118-119, and Bromwich's *Infinite Series*, p. 162, ex. 6.

Using the notation employed in the former, we have

$$u = \frac{1}{8} \{12\pi\sqrt{2}E(\sqrt{\frac{1}{2}}, \phi) - 16\pi\sqrt{2}F(\sqrt{\frac{1}{2}}, \phi)\}.$$

NUMBER THEORY AND DIOPHANTINE ANALYSIS.

188. Proposed by ELMER SCHUYLER, Brooklyn, N. Y.

Solve the congruence $3x^2 + 4x + 5 \equiv 0 \pmod{20}$.

SOLUTION BY WALTER C. EELLS, Whitworth College.

By inspection, solutions of the two congruences,

$$3x^2 + 4x + 5 \equiv 0 \pmod{4}, \quad (1)$$

$$3x^2 + 4x + 5 \equiv 0 \pmod{5}, \quad (2)$$

are $x = \pm 1$ for (1), and $x = 0, 2$ for (2).

Then we have the four systems of linear congruences,

$$\begin{array}{ll} (3) \begin{cases} x \equiv 1 \pmod{4}, \\ x \equiv 0 \pmod{5}, \end{cases} & (4) \begin{cases} x \equiv 1 \pmod{4}, \\ x \equiv 2 \pmod{5}, \end{cases} \\ (5) \begin{cases} x \equiv -1 \pmod{4}, \\ x \equiv 0 \pmod{5}, \end{cases} & (6) \begin{cases} x \equiv -1 \pmod{4}, \\ x \equiv 2 \pmod{5}. \end{cases} \end{array}$$

To solve system (3), substitute $x = 1 + 4y$ from first congruence in second congruence,

$$1 + 4y \equiv 0 \pmod{5};$$

whence

$$y = 1 + 5k,$$

and

$$x = 1 + 4y = 1 + 4(1 + 5k) = 5 + 20k.$$

Hence, $x \equiv 5 \pmod{20}$, yielding one solution of given congruence. Similarly, systems (4), (5), (6) yield $x = 17, 15, 7$, giving altogether four independent roots of the congruence $3x^2 + 4x + 5 \equiv 0 \pmod{20}$.

Also solved by H. C. FEEMSTER, LOUIS CLARK, S. LEFSCHETZ, E. B. ESCOTT.

No solution of 189 has been received.